

# **ACL** Injury

#### **Advanced Programme**

Aim to perform this programme a minimum of once per day unless prescribed otherwise. As with any new exercise, start slowly (repetitions as able) and build up as you are able within the guidelines below.

Pain should not exceed 4/10 whilst completing this exercise programme.



#### 1. Split squat

15 reps / 2-3 sets / 1 minute rest

- Place one foot forward and the other back to form a comfortable stride with the feet hip-width apart – you can gently hold on to something for balance.
- Keep the chest up and core engaged whilst simultaneously bending both knees so that the hips lower directly downwards - emphasis should be placed on pushing up through the front foot. and keeping the knee in line with the hip and ankle.
- With the trailing knee just of the floor, hold this position for a count of 3 seconds before pushing down through the feet to straighten the knees and return to the starting position.
- To progress, hold a small amount of weight in the opposite hand to the leading leg and gradually increase this as strength develops.



### 2. Single leg deadlift

15 reps / 2-3 sets / 1 minute rest

- Perform this exercise standing on the affected leg and have a wall/stable object nearby to assist with balance.
- Shift your weight on to the affected leg and 'hover' the other leg off the floor.
- Bend forward from the hip to move towards touching your toes.
- At the same time, the trailing leg should stay straight and will travel backwards to act as a counterbalance whilst keeping your spine straight
- As you get close to touching your toes, you should feel a pull in the hamstrings and buttock of the standing leg.
- Squeeze the buttock muscles to reverse the movement and return to standing.
- To progress, hold a small amount of weight in the opposite hand to the leading leg and gradually increase this as strength develops.

30 reps / 2-3 sets / 1 minute rest



## 3. Weighted squat

- Stand upright with your feet hip-width apart and a weight placed evenly across your shoulders (back squat) or holding it securely against your chest (goblet squat).
- Engage the abdominal region and in a controlled manner, sit back as if you are sitting into a chair.
- At the same time, your head and chest will come forwards to maintain your balance, aim to keep your back straight.
- Go down as far as you feel comfortable or until your thighs are parallel with the floor.
- Come back up to standing and repeat.

We recommend consulting a musculoskeletal physiotherapist to ensure exercises are best suited to your recovery. If you are carrying out an exercise regime without consulting a healthcare professional, you do so at your own risk. If you have any concerns whilst completing these exercises, please contact a healthcare professional.