# Ankylosing Spondylitis

Intermediate Programme

Aim to perform this programme a minimum of once per day unless prescribed otherwise. As with any new exercise, start slowly (repetitions as able) and build up as you are able within the guidelines below.
Pain should not exceed 4/10 whilst completing this exercise programme.

10 reps / 1-2 sets / 1 minute rest


1. Forward bending in standing

- Start standing upright.
- Place your hands on your thighs.
- Slide your hands down your legs allowing you back to bend as you do.
- Go down as far as you feel comfortable and then slowly return to standing.

15 reps / 2-3 sets / 1 minute rest

2. Bridge with leg lift

- Lie on your back on the floor or bed with your knees bent at 90 degrees and your feet on the floor and your arms down by your side.
- Push your heels down into the floor to gently lift your hips upwards. Lift as high as you feel comfortable but no higher than a straight line between hips, knees and shoulders.
- If you feel discomfort, then try squeezing your tummy and bum muscles before starting the lift.
- From this position lift one leg slightly off the floor and return.
- Repeat on the opposite side.
- Try and keep the pelvis still.

15 reps / 2-3 sets / 1 minute rest

3. 4-point kneeling superman

- Start on your hands and knees.
- Make sure that your back is straight and that your hands are under your shoulders and knees under hips.
- From this position straighten one leg out behind you with the foot slightly off the floor.
- Return back to the start position and repeat with the other leg.
- If this feels easy then you can add the arm. Do this by straightening the opposite arm in front at the same time as the leg.

We recommend consulting a musculoskeletal physiotherapist to ensure exercises are best suited to your recovery. If you are carrying out an exercise regime without consulting a healthcare professional, you do so at your own risk. If you have any concerns whilst completing these exercises, please contact a healthcare professional.